divided into five rounded, unequal lobes. Stamens inserted in the tube of the corolla, hairy, with oblong anthers. Germ inferior, roundish; style longer than the corolla; stigma peltate. The fruit is an oval berry of a deep orange yellow,* hairy, somewhat three sided, crowned with the calyx, containing three cells and three hard, bony, furrowed seeds, from which the name of the genus is taken.

This plant was made the subject of an interesting communication to the Linnæan society of New England, by Dr. John Randall. The experiments made by him on its medical uses and pharmaceutical preparations were numerous, and serve to throw much light on its properties. In trying the solvent powers of water and alcohol, he found that water afforded a much greater quantity of extract than alcohol, and that the spirituous extract was perfectly soluble in water, whence he infers that no resin in a pure state exists in the plant. He discovered no volatile oil by distillation, nor any other principle of activity in water distilled from the plant. He concludes also, that

* Pursh observes that the flowers and berries are *purple*. In all the specimens I have examined, which have not been few in number, the fruit was of a bright orange colour. If Pursh has seen a plant with purple berries, it is probably a different species from the true plant of Linnæus and Dillenius, which had "*fructus lutescentes*."

no free acid exists in this vegetable. Of the different parts submitted to examination, the leaves yielded the greatest quantity of soluble matter, but the root afforded that of the greatest activity. By decoction and evaporation with water an ounce of the dried stalks afforded one drachm of extract; an ounce of the dry roots, two drachms and two scruples, and the same quantity of leaves half an ounce. From a similar treatment of equal portions with alcohol, rather more than half the above quantities of extract were obtained.

The sensible qualities of the root were found essentially different from those of the herb. Both of them possess a large share of bitterness, but the root has also a nauseous taste and smell, somewhat approaching to those of ipecacuanha. The medical properties of the Triosteum are those of an emetic and cathartic. In the above dissertation, about thirty cases are detailed, in which different preparations and quantities of the article were given to various persons with a view to their medicinal effects. The general inference to be made from them is, that the bark of the root acts with tolerable certainty as an evacuant upon the alimentary canal, both by emesis and catharsis. When given alone, either in powder or decoction, the instances of its failure were not many, and

JACOB BIGELOW *American Medical Botany* BOSTON, 1817-1820 CARY GRAPHIC ARTS COLLECTION, ROCHESTER INSTITUTE OF TECHNOLOGY Octavo